

INVITED REVIEW ARTICLE

Kidney organoids: a pioneering model for kidney diseases

MURAT TEKGUC, RONALD C. VAN GAAL, SEBASTIEN G.M. UZEL, NAVIN GUPTA, LEONARDO V. RIELLA, JENNIFER A. LEWIS, and RYUJI MORIZANE

BOSTON, AND CAMBRIDGE, MASSACHUSETTS

The kidney is a vital organ that regulates the bodily fluid and electrolyte homeostasis via tailored urinary excretion. Kidney injuries that cause severe or progressive chronic kidney disease have driven the growing population of patients with end-stage kidney disease, leading to substantial patient morbidity and mortality. This irreversible kidney damage has also created a huge socioeconomical burden on the healthcare system, highlighting the need for novel translational research models for progressive kidney diseases. Conventional research methods such as *in vitro* 2D cell culture or animal models do not fully recapitulate complex human kidney diseases. By contrast, directed differentiation of human induced pluripotent stem cells enables *in vitro* generation of patient-specific 3D kidney organoids, which can be used to model acute or chronic forms of hereditary, developmental, and metabolic kidney diseases. Furthermore, when combined with biofabrication techniques, organoids can be used as building blocks to construct vascularized kidney tissues mimicking their *in vivo* counterpart. By applying gene editing technology, organoid building blocks may be modified to minimize the process of immune rejection in kidney transplant recipients. In the foreseeable future, the universal kidney organoids derived from HLA-edited/deleted induced pluripotent stem cell (iPSC) lines may enable the supply of bioengineered organotypic kidney structures that are immune-compatible for the majority of the world population. Here, we summarize recent advances in kidney organoid research coupled with novel technologies such as organoids-on-chip and biofabrication of 3D kidney tissues providing convenient platforms for high-throughput drug screening, disease modelling, and therapeutic applications. (Translational Research 2022; 000:1 – 17)

Abbreviations: iPSC = induced pluripotent stem cell; CKD = chronic kidney disease; ESKD = end-stage kidney disease; IM = intermediate mesoderm; MM = metanephric mesenchyme; UB = ureteric bud; NPCs = nephron progenitor cells; MET = mesenchymal-to-epithelial transition; SPCs = stromal progenitor cells; EPCs = endothelial progenitor cells; HDR = homology-directed repair; APCs = antigen-presenting cells; PKD = polycystic kidney disease; GBM = glomerular basement

From the Nephrology Division, Department of Medicine, Massachusetts General Hospital, Boston, Massachusetts; Harvard Medical School, Boston, Massachusetts; Harvard Stem Cell Institute (HSCI), Cambridge, Massachusetts; Wyss Institute for Biologically Inspired Engineering, Harvard University, Boston, Massachusetts; School of Engineering and Applied Sciences, Harvard University, Cambridge, Massachusetts; Center for Transplantation Sciences, Department of Surgery, Massachusetts General Hospital, Boston, Massachusetts.

Submitted for Publication April 11, 2022; revision submitted June 14, 2022; Accepted for Publication June 16, 2022.

Reprint requests: Ryuji Morizane, Nephrology Division, Department of Medicine, Massachusetts General Hospital, Boston, MA. E-mail address: rmorizane@mgh.harvard.edu.

1931-5244/\$ - see front matter

© 2022 Elsevier Inc. All rights reserved.

<https://doi.org/10.1016/j.trsl.2022.06.012>

membrane; CNS = congenital nephrotic syndrome; PEC = parietal epithelial cell; NPs = nephron progenitors; TMA = thrombotic microangiopathy; HLA = human leukocyte antigen;

INTRODUCTION

The kidney is responsible for the regulation of fluid, electrolyte, and pH balance in the human body via plasma filtration, reabsorption of essential nutrients, and secretion of the metabolic waste.⁹⁴ The kidney is composed of more than 20 distinct cell types, that are necessary for its function to maintain homeostasis. While tubular cells have the capacity to self-repair after injury, repeated and/or severe renal injury cause irreversible damage to the kidney, leading to its dysfunction.^{21,94} Chronic kidney disease (CKD) constitutes a major global healthcare problem causing a heavy financial burden with its high prevalence in 13.4% of the global population and 35% of the people over 70 years.^{44,53} In addition, CKD is an independent risk factor for cardiovascular diseases.⁵³ CKD is directly or indirectly caused by many types of disorders including hereditary diseases, metabolic syndrome and chronic autoimmune and/or inflammatory diseases or as a cumulative result of drug-induced nephrotoxicity.⁴⁶ Loss of functional units of the kidney causes progressive loss of renal function over time due to its limited regeneration capacity, leading to end-stage kidney disease (ESKD), which requires patients to initiate renal replacement therapy.

There have been many efforts to create novel sources for kidney regeneration during the past decade. One approach is to derive kidney cells and tissues from human pluripotent stem cells (hPSCs). Detailed analyses of the embryonic origins of kidney lineages have enabled researchers to induce its differentiation process using pluripotent stem cells *in vitro*.^{75,92} By mimicking signal cascades and growth factor stimulation of embryonic kidney development, directed differentiation protocols have been developed to generate kidney organoids that consist of segmented-nephrons and stromal cells.^{45,68} These kidney organoids can now serve as a new tool for patient-specific drug screening and disease modeling. Coupled with CRISPR/Cas9 genome editing, genetic kidney disease models are established with their isogenic control organoids, complementing animal experimentation.³² Towards the ultimate goals of kidney regeneration, significant advances have been made in biomanufacturing organ-specific tissues at scale to overcome the challenge of large tissue fabrication *in vitro*.⁸⁹ Moreover, the generation of hypoinmunogenic hPSCs by genome editing may mitigate the immune rejection against the bioengineered organs. In this review, we summarize the recent advances in kidney organoids, disease models, and efforts that are underway to build

transplantable kidneys that might revolutionize the care of patients with ESKD.

Kidney structure and development. Nephrons are the functional unit of the kidney, responsible for the production and excretion of urine via blood filtration and active transport. Each nephron consists of glomerular and tubular segments. Glomerular podocytes form filtration units with capillary networks integrated within the Bowman's capsule, filtering blood to produce ultrafiltrate as primitive urine. Glomeruli are structurally continuous with the proximal tubules, loops of Henle, distal tubules, and connecting tubules and/or collecting ducts⁴⁵ (Fig 1). After filtration of the renal blood from the glomerular capillaries through their highly fenestrated endothelium,⁶⁰ this glomerular ultrafiltrate flows from Bowman's space to the tubular lumens. Tubular cells are surrounded by peritubular capillaries, and 99% of the glomerular ultrafiltrate is reabsorbed from the tubular lumens to peritubular capillaries. The remaining portion is further subject to tubular secretion of ions and organic solutes, determining the final composition of urine.^{43,84} Resorptive and secretory processes are facilitated by active transport in the tubular epithelial cells together with the peritubular capillary endothelia. Finally, urine produced by nephrons is collected to the bladder through the collecting duct and ureter.

Kidneys developmentally originate from the intermediate mesoderm (IM). During the formation of the mammalian kidney, three distinct renal rudiments namely pronephros, mesonephros, and metanephros arise from anterior to posterior areas in the IM.⁴⁵ Pronephros, mesonephros, and the ureteric bud (UB) are derived from the anterior IM (aIM) whereas metanephric mesenchyme (MM) is derived from the posterior IM (pIM) (Fig 2).^{29,43} While the pronephros and mesonephros spontaneously degenerate as non-functional structures, the Wolffian duct persists as the collecting duct of the metanephric kidneys.^{34,86} Although the previous literature pointed out the *Osr1*⁺ metanephric progenitors in the IM as the common ancestry of the MM and UB,⁷⁶ the subsequent lineage-tracing mouse studies demonstrated the distinct origins of UB and MM lineages. The UB, the caudal outpouching of the Wolffian duct, arises from *Pax2*⁺*Osr1*⁺ progenitors in the anterior IM while the MM is derived from *Pax2*[−]*Osr1*⁺ progenitors in the posterior IM, that latter further differentiating into *Osr1*⁺*Six2*⁺*Sall1*⁺*WT1*⁺ nephron progenitor cells (NPCs).⁹² NPCs give rise to segmented nephron structures through reciprocal induction with the UB.^{45,76} When invaded by the UB, the MM




Fig 1. The structure of a mature nephron unit. The upper schema designates each unit of a nephron with a different function. The glomerulus, composed of Bowman's capsule and the glomerular capillary network, is responsible for the filtration of glomerular blood flow. The glomerular capillary network, formed by the afferent arteriole, is assembled into the efferent arteriole. The lower illustration shows the magnified microstructure of the glomerular filtration barrier, which is composed of glomerular fenestrated endothelial cells, glomerular basement membrane, and podocytes (visceral epithelial cells). The secondary foot processes of podocytes are attached to each other by intervening filtration slits forming the slit diaphragm. On the other hand, Bowman space is lined by the parietal epithelial cell layer. The ultrafiltrate, drained from the Bowman space into the tubules (proximal tubule, loops of Henle, and distal tubule), is further processed into the urinary output by absorption and secretion. The urinary drainage is finally transmitted into the collecting duct via the connecting tubule.

supports the extensive branching of UB tips through the secretion of glial cell-derived neurotrophic factor (GDNF), while the cap mesenchyme surrounding the UB tips undergoes a mesenchymal-to-epithelial transition (MET) via UB-derived Wnt signal activation.^{24,29,111} The pIM-derived MM is composed of the heterogeneous cell populations of NPCs, stromal progenitor cells (SPCs), and endothelial progenitor cells (EPCs), which give rise to the nephron structures and their surrounding interstitial tissue (Fig 2).

aIM-derived UB differentiates into the collecting duct network which is responsible for draining the urine to the bladder via ureters.⁴⁵

Differentiation of kidney organoid from hPSCs. Based on the knowledge of mammalian kidney development, protocols have been established to generate nephron kidney organoids from hPSCs.^{43,46} These protocols are based on the approach of “directed differentiation” whereby sequential treatment of growth factors is developed to follow the developmental stages of the

Embryological development of kidney




Fig 2. The embryological development of kidney at different stages. The upper schema shows the formation of paraxial, intermediate, and lateral plate mesodermal layers, derived from the primitive streak as a result of cellular immigration along its craniocaudal axis. The lower illustration reveals the embryological stages of pronephros, mesonephros, and metanephros. Pronephros, mesonephros, and the ureteric bud (UB) are originated from the anterior intermediate mesoderm (aIM) whereas metanephric mesenchyme (MM) is derived from the posterior intermediate mesoderm (pIM). The lower right schema reflects the microstructure of MM- and UB-derived progenitor cells.

kidney. Three-dimensional (3D) culture technologies enabled embryonic and adult stem cells to differentiate into kidney tissues by potentiating their self-organizing capacity to form multi-cellular tissues. Organoids

derived from hPSCs serve as a novel tool for disease studies. In addition, organoids produced in 3D culture can be used as organ building blocks as they have similar basic structural and functional features exhibited by

their native counterparts in the human body.^{20,43} In this section, we summarize the differentiation protocols of kidney organoids.

Taguchi et al. achieved a milestone in kidney developmental research by elucidating the distinct origins of ureteric and metanephric lineages.⁹² Nephron progenitor cells from both mouse embryonic stem cells (ESCs) and human induced pluripotent stem cells (iPSCs) were generated by directing their differentiation towards pIM-derived MM cells, which subsequently commenced their maturation into nephron epithelia when cocultured with mouse embryonic spinal cords. First, OCT3/4⁺ hPSCs were treated with BMP4 followed by activin A and FGF2 to shift their pluripotent status to the epiblast stage. Then, the differentiation was initiated by BMP4 and a glycogen synthase kinase 3 (GSK-3) inhibitor, CHIR99021 (CHIR) to induce T⁺CDX2⁺TBX6⁺HOX11⁺ posterior nascent mesoderm.⁷ The subsequent treatment of activin A, BMP-4, retinoic acid, and a moderate dose of CHIR progressed the differentiation into OSR1⁺WT1⁺HOX11⁺ pIM cells. They then applied FGF9 and a low dose of CHIR to induce SIX2⁺PAX2⁺SALL1⁺ MM cells. Finally, these MM cells were co-cultured with mouse embryonic spinal cords to induce tubular and podocyte epithelial cells.

Morizane et al. developed a rapid and efficient protocol with defined components, which generated SIX2⁺SALL1⁺WT1⁺PAX2⁺ NPCs with ~90% efficiency from both human embryonic stem cells and iPSCs within 8 to 9 days (Fig 3).⁷⁵ Furthermore, these NPCs were converted into PAX8⁺LHX1⁺ renal vesicles, which had the potential to self-organize into their *in vivo* counterparts; nephron structures including NPHS1⁺PODXL⁺ glomerular podocytes, LTL⁺CDH2⁺ proximal tubules, CDH1⁺UMOD (Uromodulin)⁺ loops of Henle, and CDH1⁺BRN1⁺ distal tubules in an organized and continuous manner.^{74,75} The differentiation protocol was arranged according to two findings that the egression timing of mesodermal progenitors from the primitive streak determines the anterior-posterior patterning of the mesoderm,²⁷ and the mesodermal precursors from the posterior primitive streak transform into lateral plate mesoderm instead of intermediate mesoderm.⁵⁸ Posterior mesoderm progenitors leave the primitive streak later than the precursor cells of the anterior mesoderm indicating that the posterior intermediate mesoderm cells originate from the late-stage primitive streak. Based on this embryologic specification, high-dose and long-term CHIR treatment was applied together with the BMP4 inhibitor that suppressed the differentiation of lateral plate mesoderm components. During *in vivo* nephrogenesis, Activin A (TGF- β superfamily member) activates Hoxd11

expression, which stimulates the formation of pIM-derived metanephros.^{38,45,71} Thus, after the removal of CHIR, activin A treatment was applied between days 4 and 7 to induce OSR1⁺WT1⁺HOXD11⁺ pIM cells which lacked aIM markers, PAX2 and LHX1.^{75,92} Next, SIX2⁺SALL1⁺WT1⁺PAX2⁺ NPCs were generated when pIM cells were treated with FGF9 for 1 or 2 days. Previous studies have shown that UB-derived Fgf9 and Fgf20 are necessary to maintain the stemness and survival of cap mesenchyme in mice,⁶ while UB-derived Wnt9b signaling supports the mesenchymal-epithelial transition of MM to nephron epithelial cells.¹⁷ In addition, the induction of Fgf8 and Fgf9 by Wnt9b signaling was also reported,^{16,63} referring to the interaction between Fgf9 and Wnt signaling during the reciprocal induction of MM and UB. To mimic the MET in murine NPCs as a result of this reciprocal induction by the UB-derived Wnt9b and Fgf9 signaling,¹⁷ hPSC-derived NPCs were further exposed to a low-dose CHIR pulse together with the ongoing FGF9 following their transfer to 3D suspension culture and brief centrifugation in the ultra-low-attachment plates.⁴⁵ After the CHIR pulse, NPCs were transformed to PAX8⁺LHX1⁺LAM⁺ renal vesicles by day 14 of the differentiation, which is consistent with the previous finding that Wnt signaling is abrogated by Dkk1, a key downstream effector of *Lim1* (*Lhx1*) gene during the renal vesicle formation.⁸¹ Finally, these renal vesicles spontaneously completed their MET in the absence of any additional factors by forming elongated epithelial nephron structures composed of NPHS1⁺PODXL⁺ glomerular podocytes, LTL⁺AQP1 (Aquaporin-1)⁺ proximal tubules, CDH1⁺AQP1⁺ descending loops of Henle, CDH1⁺UMOD⁺ thick ascending loops of Henle, CDH1⁺UMOD⁻ distal tubules, and CDH1⁺AQP2 (Aquaporin-2)⁺ connecting tubules.^{74,75} The common main points of the differentiation protocols by Taguchi and Morizane included the induction of the late primitive streak or posterior nascent mesoderm with long-term WNT activation by CHIR and modified BMP4 signaling (further activation of BMP4 by Taguchi protocol, suppression of excessive BMP4 signaling by Morizane protocol), the differentiation of late primitive streak to pIM with Activin A, the maturation of MM by FGF-9, and the induction of MET in NPCs through transient activation of WNT signaling by either pulse CHIR treatment (Morizane) or coculture with embryonic spinal cords (Taguchi).^{75,92}

Although most researchers have performed suspension cell culture for organoid differentiation in low-attachment wells, Freedman et al. used a Matrigel sandwich method to generate kidney organoids in the 3D microenvironment via cavitated epiblast formation from hPSCs. Before starting directed differentiation,

In vitro differentiation of nephron organoids




Fig 3. Directed differentiation of kidney organoids by the Morizane protocol. The timescale of this protocol indicates the temporal stages of the organoid development and the temporal supplementation of the differentiation factors. The lower left image (scale bar, 1 mm) by confocal microscopy shows DAPI⁺SIX2⁺ nephron progenitor cells (NPCs) at Day 9 while the lower right image (scale bar, 100 μ m) clarifies the development of DAPI⁺Podocalyxin⁺LTL⁺CDH1⁺ nephron organoids at Day 21. PS: primitive streak. IM: intermediate mesoderm. MM: metanephric mesenchyme.

hPSCs were sandwiched between two layers of Matrigel, forming epiblast spheroids that contained a hollow lumen surrounded by monolayer epithelia. These epiblast spheroids were first treated with CHIR for 1.5 days and subsequently exposed to B27-supplemented media by a two-step protocol. After day 10, spheroids were transformed into convoluted tubular organoids which contained PODXL⁺WT1⁺SYNPO⁺ podocyte-like cell aggregates, LTL⁺ proximal and CDH1⁺ distal tubular structures, and CD31⁺vWF⁺ endothelial compartments.³³

Takasato et al. was first to report the synchronous induction of the nephrons and collecting ducts from

aIM progenitors, together forming a self-organizing kidney structure.⁹³ They modified the duration of initial CHIR treatment to co-induce both nephron and collecting duct lineages.⁹⁴ By using CHIR for 4 days, the optimal duration for the induction of MM together with UB, and subsequent application of FGF-9 for 3 days, they simultaneously induced MM- and UB-derived components including WT1⁺NPHS1⁺ glomeruli, LTL⁺CDH1⁻ early proximal tubule (day 11), LTL⁺CDH1⁺ maturing proximal tubule (day 18), GATA3⁻LTL⁻CDH1⁺ early distal tubule, GATA3⁺CDH1⁺ collecting duct.⁹⁴ Although GATA3 was used to distinguish between distal tubules and collecting

ducts, later studies indicated that GATA3 can also be expressed by the distal tubules and connecting segments of the nephrons,²² suggesting the necessity of different protocols to specifically induce UB lineages. Additionally, the same group reported that human kidney organoid-derived glomeruli (OrgGloms) cultured in 3D exhibited proper apicobasal polarization and high expression of podocyte-specific genes such as *NPHS1*, *NPHS2*, and *PODXL* in addition to their rich content of glomerular basement membrane matrices when compared to conditionally immortalized human podocytes cultured in 2D.^{48,94} These polarized podocytes of OrgGloms expressed *PODXL* at the apical surface and *NPHS1* and *NEPH1* proteins on the basal side which are critical for the filtration function of the kidney.^{56,90} These pioneering studies of kidney organoid differentiation opened new avenues to investigate kidney development and diseases, followed by numerous studies that utilized kidney organoids for disease models.

Disease models using kidney organoids. There are many challenges in using two-dimensional (2D) cell cultures and animal models for drug screening and disease modeling.⁴³ Conventional monolayer cell cultures lack the tissue complexity and 3D biochemical and biomechanical microenvironments present in human organs.³⁰ Although murine models are widely used to explore the pathophysiological mechanisms of the genetic or autoimmune and/or inflammatory diseases for the development of novel therapeutic agents prior to clinical studies in patients, there has been a growing concern for the translation of animal studies to humans. The published literature, which compared the differential gene expression profiles of mice and humans via various microarrays during the acute inflammatory responses, has shown that even though different etiologies of acute inflammatory disorders produce very similar genomic responses among different patients, these gene expression signatures in the patients are not correlated with the corresponding profiles in murine models.⁶¹ Furthermore, there have been growing ethical concerns for animal experiments due to the importance of animal welfare and rights, gradually limiting animal experimentation.² Therefore, human kidney organoids are of great interest for disease modeling and drug screening owing to their potential to minimize these animal studies.

(1) Kidney organoid models for tubular injury and fibrosis

Drug-induced nephrotoxicity, frequently documented among patients in the intensive care unit (ICU), has remained challenging during ongoing treatment for their primary diseases, causing acute kidney

injury (AKI) and CKD progression.¹⁰⁰ For this critical problem, human kidney organoids were tested for nephrotoxicity assays using gentamycin and cisplatin.⁷⁵ Gentamycin is a commonly used antibiotic which causes proximal tubular injury, while cisplatin, a chemotherapeutic agent used for a wide range of cancers, also induces dose-dependent proximal tubular damage.^{75,105} Both drugs particularly damaged the proximal tubules of organoids as evidenced by the expression of kidney injury molecule-1 (KIM-1) on the luminal surface of LTL⁺ proximal tubules.¹⁰² Interestingly, podocytes did not exhibit injury phenotypes as suggested by the absence of a DNA damage marker, γ H2AX, when low doses of these drugs were used. This early work suggested that kidney organoids express drug transporters in a segment-specific manner whereby cell type-specific responses to varied drugs can be tested in the organoid model.

Gupta et al. recently developed an organoid model of kidney injury and repair by applying repetitive cisplatin exposure. This study demonstrated that kidney organoids arose from co-induced nephron and stromal progenitor cells, which led to the generation of segmented nephrons admixed with stromal cells. Given the presence of discernible kidney compartments, the authors histologically monitored intrinsic repair (reversible injury) vs incomplete repair (irreversible injury) throughout a protocol of repeated cisplatin exposure, analogous to cycles of chemotherapy.^{44,88} Following an initial 24 hours exposure to low-dose cisplatin, proximal tubules specifically manifest injury markers of DNA damage and KIM1, as previously reported.⁷⁵ Knowing that injured tubular epithelial cells compensate by proliferative expansion to preserve kidney architecture during intrinsic repair,⁵⁷ the authors demonstrated the induction of intrinsic repair phenotypes of proliferation and dedifferentiation, as evidenced by the reactivation of kidney developmental genes, *SOX9* and *PAX2*. Meanwhile, the DNA damage response triggered the elevation of homology-directed repair (HDR) protein, Fanconi anemia complementation group D2 (FANCD2). In a longitudinal study after the initial injury, the resolution of DNA damage by FANCD2 activation during the intrinsic repair phase terminated the proliferative phenotype with preservation of tubular structures reflecting the reversibility of the injury consistent with completed intrinsic repair. However, when the acute cisplatin injury was repeated biweekly for 5 treatments, incomplete repair phenotypes of tubular atrophy, myofibroblast activation, and interstitial fibrosis signified irreversible injury.⁴⁴ Loss of FANCD2 in DNA damaged tubules correlated with the transition from intrinsic to incomplete repair in kidney organoid samples. Mammalian single cell

transcriptomic datasets of ischemic, obstructive, and immune-mediated kidney injury were interrogated to confirm the generalizability of tubular FANCD2 correlating with intrinsic repair, whereas human fibrotic biopsy samples suggested that tubular FANCD2 inversely correlated with incomplete repair. To extend beyond a correlation, a small molecule inhibitor that upregulates HDR was found to rescue FANCD2 in injured tubules, partially reversing tubular atrophy, myofibroblast activation, and interstitial fibrosis.⁴⁴ It was concluded that HDR is a mechanism underpinning intrinsic repair following tubular injury in this organoid model developed in the absence of the potential pro and anti-fibrotic effects of the immune system. Of note, the single cell analysis in this study suggested the myofibroblast transition from mesenchymal fibroblast populations in kidney organoids, which is consistent with a recent study in human kidney samples.⁶⁴ In addition, another hierarchical single-cell transcriptional profiling model, generated through the multiple single cell RNA-seq datasets of human fetal kidney (HFK), has reclassified the component cellular identities, such as stromal and NPC-like cells within the published kidney organoid datasets.¹⁰⁸ As the HFK dataset grouped stromal cells into 4 subpopulations such as stromal progenitor cells (SPC), cortical stroma (CS), medullary stroma (MS), and mesangial cells (MesS), the organoid stroma was mainly composed of cortical stromal cells that are the origin of myofibroblasts.

(2) Kidney organoids under the inflammatory condition

During acute kidney injury, repair, and fibrosis, immune and/or inflammatory responses play an important role in modulating intrinsic and incomplete repair. Lemos et al. used kidney organoids as a fibrosis model by exposing them to an inflammatory cytokine, IL-1 β .⁶⁷ Kidney organoids treated with IL-1 β developed tubulointerstitial fibrosis with α SMA⁺ myofibroblast activation and collagen deposition. IL-1 β signaling through the IL-1 receptor-associated kinase 4 (IRAK4) promoted the upregulation of *MYC* downstream target genes, containing cell cycle regulators and glycolytic-proliferative transcriptional signature, which eventually led to the proliferation of kidney stromal cells (SCs) and their transdifferentiation into myofibroblasts, and the deposition of fibrotic matrices in kidney organoids.

The investigation of the immune system in laboratory mouse strains is significantly limited by the minimum genetic diversity and predominance of an immature immune system that resembles neonatal immune reaction in organs such as the kidney.^{12,49} Therefore, human organoids could better reproduce the

microenvironment of the native kidney with its complex reciprocal interaction between T and B lymphocytes and antigen-presenting cells (APCs), that involves antigen presentation and simultaneous expression of costimulatory molecules to activate antigen-specific effector T and B lymphocytes during adaptive immune response.¹⁰⁹ Kidney organoids may allow us to better understand the pathophysiology of immune-mediated kidney diseases such as auto-immune glomerulonephritis and enable drug testing for novel therapeutic strategies. A recent example is the investigation of high-risk apolipoprotein-1 (APOL1) variant genes in kidney disease, which is a leading cause of kidney disease in African Americans.³⁹ Using different combinations of APOL1 variants generated by CRISPR/Cas9 genome editing, Chun et al. demonstrated the role of interferon-gamma signaling that drove lipid accumulation and podocyte toxicity in kidney organoids with high-risk APOL1 variants, which could be reversed by inhibition of diacylglycerol O-acyltransferases 2.¹⁹

The kidneys are often involved in systemic autoimmune and autoinflammatory diseases with consequent deposition of antibody, complement, and/or immune complexes in glomeruli. Several studies pointed out the contribution of non-immune cells of the kidney in such immune-mediated glomerulonephritis. Interestingly, podocytes can function as antigen-presenting cells to induce CD4⁺ T cells and also CD8⁺ T cells via cross presentation during both *in vitro* and *in vivo* settings, while proximal tubular epithelial cells activate the proliferation and cytokine production of CD4⁺ T cells when cocultured together.^{14,40} Furthermore, the presence of co-inhibitory receptors such as PD-L1 in kidney tubules suggests their potential role in actively controlling the immune response.^{47,91} The unique contribution of immune and non-immune cells to kidney injury and fibrosis in the organoid models deserves further investigation to clarify the complex interactions of these cells during immune-mediated nephritis. Further, autoimmune and/or autoinflammatory diseases that involve the interaction between epithelia and immune cells could be studied using kidney organoids in combination with genome editing technologies that can modify the genes of interests.⁵ The effects of immunosuppressive modalities such as regulatory T cells or CTLA-4 immunoglobulin could also be tested in organoid models⁹⁷ for the future applications.

(3) Kidney organoids for hereditary kidney diseases with tubulopathy

Adult-onset autosomal dominant polycystic kidney disease (PKD), the most common form of hereditary renal disorder with an incidence of approximately

1:1000 live births, is characterized by the formation of renal cysts derived from tubular epithelial cells, leading to the progressive loss of renal function in enlarged kidneys.⁵⁴ This life-threatening disorder is caused by the mutations in *PKD1* or *PKD2* genes encoding polycystin 1 (PC1) and polycystin 2 (PC2) proteins, which form a heterodimeric complex to function in primary cilia.¹⁰ Freedman et al. introduced a PKD model by using gene-edited human kidney organoids.³³ Following the generation of *PKD*-mutated hPSCs by CRISPR/Cas9 genome editing, they demonstrated the formation and expansion of cyst-like structures in the tubular organoids differentiated from *PKD1* or *PKD2*-mutants. Although ADPKD cysts arise pre-dominantly from the collecting duct, the cysts in this study were localized in the LTL⁺ tubules.³³ In 2017, they found that PC1 protein was responsible for the adhesion of tubular epithelial cell compartment to the underlying extracellular matrix (ECM) microenvironment, which was required for the maintenance of the tubular shape and integrity. In addition, PC1 participated in the remodeling of ECM microenvironment, which eventually limited the cyst formation.²⁵ Therefore, it was shown that removal of adherent ECM components further promoted the cystogenesis of PKD-mutant/PC1-deficient organoids, especially in early stages of the disease. Furthermore, cAMP signaling was shown to induce the enlargement of cyst-like structures, contributing to the pathogenesis of PKD.

Nephronophthisis (NPHP)-related ciliopathy is a progressive fibrocystic kidney disease characterized by shrunk or normal-sized fibrotic kidneys with small cysts at the corticomedullary junction in contrast to PKD presented with huge cysts.¹³ Histological findings are characterized by the disruption of the tubular basement membrane, tubular atrophy and/or dilatation, and interstitial fibrosis, eventually resulting in end-stage kidney disease.³² Little et al. demonstrated the underlying pathology of NPHP in patient kidney organoids by comparing the organoids of an identified NPHP proband (PR) patient with its isogenic control organoids derived from their gene-corrected (GC) iPSCs. By using CRISPR/Cas9 technology, the GC iPSCs were originally obtained from the same NPHP PR patient with compound-heterozygous mutations detected in the *IFT140* gene.³² When kidney organoids derived from both PR and GC iPSCs were compared, PR organoid tubules exhibited shortened and club-shaped primary cilia, whose phenotype was rescued by the gene correction. Consistent with the downregulation of genes implicated in apicobasal polarity, cell-cell junctions, and dynein motor assembly, spheroid cultures demonstrated the apicobasal polarization defect in PR organoid tubules unlike GC organoids. However, the

cystic kidney phenotype was not seen in this study, that might be due to the limitations of static organoid models. A recent study reported that cystic tubular cells derived from ADPKD patients performed basoapical fluidic pumping in the opposite direction compared to normal tubular cells, revealing the importance of hydraulic pressure in ADPKD pathophysiology.¹⁸ Therefore, novel approaches, which can simulate *in vivo* fluidic microenvironment, such as organoids-on-chip approaches can be used in the future to investigate the cyst formation in the ciliopathic organoids in the presence of fluidic shear stress.^{43,55} Nevertheless, this work showed an approach to correct patient mutations by CRISPR/Cas9 for generation of an isogenic control, which validated the ciliary pathogenesis in the PR organoids.³²

(4) Kidney organoid models for glomerulopathy

A glomerulus, a filtration segment of each nephron, comprises fenestrated endothelia, mesangial cells, podocytes, and parietal epithelial cells.^{48,87} Podocytes, anatomically defined as glomerular visceral epithelial cells, form the glomerular filtration barrier with their foot processes (FPs) and the capillary endothelial cells.^{31,48,87} FPs create interdigitating structures with the adjacent podocytes, forming slit diaphragms that prevent the leakage of plasma proteins from the circulating blood together with the glomerular basement membrane (GBM) (Fig 1).^{28,31,42,48} Podocyte injury and its structural alterations (eg, foot process effacement) cause nephrotic syndrome that manifests massive proteinuria, severe edema, dyslipidemia, leading to progressive loss of kidney function.^{31,48,106} In 2018, Little et al. isolated organoid-derived glomeruli (OrgGloms) expressing *mTagBFP2* fluorescent reporter gene in their developing podocytes, and performed toxicity screening using doxorubicin (Adriamycin), a chemotherapy medication, at various concentrations.⁴⁸ Doxorubicin induced loss of BFP2 signal in OrgGloms, the fragmentation and destruction of podocytes in a dose dependent manner.

The podocyte unique structures are supported by slit diaphragm proteins including nephrin, podocin, and synaptopodin.³¹ Wilms tumor-1 (*WT1*), earliest podocyte marker during kidney development, is identified as one of the upstream transcription factors which regulate these podocyte genes.^{28,75} To reveal the epigenetic transcriptional reprogramming by *WT1*, Ettou et al. performed genome-wide analysis of *WT1* DNA binding and evaluated podocyte gene expression during the course of Adriamycin (ADR)-mediated injury in both murine models and human organoids. They demonstrated that *WT1* is responsible for the maintenance of open chromatin at podocyte genes for their active

transcription.³¹ Although ADR-induced injury initially promoted an enhanced open chromatin state at *WT1* target genes such as *NPHS2* (Podocin) and *SYNPO* (Synaptopodin), the chromatin status was subsequently converted from active to repressive, downregulating podocyte genes. Hence, this study suggested that *WT1* is the key transcription factor for podocyte survival during injury and the transcriptional reprogramming of podocyte gene expression can be a novel therapeutic target for CKD.³¹ These two studies demonstrate that 3D human kidney organoids can be utilized to study podocyte injury induced by toxicants.

Notably, Little et al. also showed the application of kidney organoids to model the congenital nephrotic syndrome (CNS) caused by the mutations in *NPHS1* (nephrin) and *NPHS2* (podocin).^{48,99} Scanning electron microscopy findings of the OrgGloms derived from the CNS patient carrying *NPHS1* gene mutations pointed out the large hypertrophied podocyte bodies, while quantitative western blot analysis confirmed the significant reduction in both nephrin and podocin proteins in the patient derived-OrgGloms compared to the control group. A similar study using hiPSC-derived kidney organoids from a patient with *NPHS1* missense mutation demonstrated the formation of impaired slit diaphragm (SD) in the organoid podocytes.⁹⁵ These *NPHS1* mutant podocytes of the organoids were deprived of “pre-SD domain” structure, which was defined as the transit state of rod-like Nephrin protein shifting between the lateral and basal sides of podocytes. Genetic correction of this point mutation restored the nephrin expression, its phosphorylation, and the formation of lateral and basal nephrin⁺ pre-SD domains resulting in the generation of proper SD.⁹⁵ Both of these organoid studies modeling CNS revealed different aspects of the structural disruption caused by the nephrin mutation.

One study focused on another cell type of glomeruli, parietal epithelial cells (PEC) (Fig 1). Nishinakamura et al. demonstrated PAX2 is indispensable for morphological maturation of PEC using kidney organoids and transcription activator-like effector nucleases (TALENs). Interestingly, PAX2 was found to be dispensable for *in vitro* nephron formation from human iPSCs.⁶² Not only human nephron progenitors (NPs) were formed from PAX2-depleted hiPSCs, but, also they were able to become epithelialized to form glomerular and renal tubular structures. However, glomerular PECs remained to be columnar epithelia, lacking the ability to transform into the squamous morphology. Hence, they confirmed that human PAX2 protein is not required for the mesenchymal-to-epithelial transition of nephron progenitors to develop normal glomeruli

and renal tubules, but indispensable for the normal differentiation of PECs.⁶²

Research work discussed above demonstrate the utility of human kidney organoids for a variety of genetic kidney diseases (ADPKD, nephronophthisis-related ciliopathies, congenital nephrotic syndrome), developmental kidney disorders (PAX2 deficiency), drug-induced glomerular or tubular toxicity, and inflammatory cytokine-mediated renal injury. Patient-derived organoids can also be used for personalized drug screening to assess the therapeutic efficacy and side effects as preclinical trials. Although kidney organoids exhibit most of the structural elements of native kidney *in vitro*,^{74,75} the formation of vascular network is limited in the static culture of organoids.^{55,75} Organoids generated by Freedman et al. contained CD31 and von Willebrand factor expressing endothelial cords while organoids by Takasato et al. developed CD31⁺KDR⁺SOX17⁺ endothelial network.^{33,94} However, these studies have not observed mature vascularized glomeruli, yet. Only after subcapsular transplantation of the organoids into mice, host-derived vascularization induced glomerular and peritubular neo-vasculogenesis.⁹ Development of vascularized kidney organoids would improve the organoid disease models for translational studies.

Optimizing kidney organoid models. Kidney organoids offer a novel platform for drug screening, disease modeling, and regenerative tissue engineering, since they contain key cell types that self-assemble into the desired nephron-like subunit microarchitectures.^{75,93} However, kidney organoids differentiated under static culture conditions lack the microvasculature needed to facilitate both filtration and reabsorption *in vitro*.^{55,66} Recently,⁵⁵ Homan and Gupta *et al.* demonstrated that subjecting kidney organoids to flow during their differentiation leads to a substantial upregulation in their endothelial progenitor population, which promotes the formation of a pervasive microvascular network and enhances organoid maturation. Specifically, under flow, the vascular invasion of podocyte-rich clusters is observed along with primitive foot process formation and an upregulation in key glomerular markers.⁵⁵ Concomitantly, vascular wrapping of proximal tubular segments is enhanced under flow that leads to improved polarity, ciliation, and upregulation of key proximal tubule markers.⁵⁵ The importance of flow during kidney organoid differentiation was also highlighted by a recent study from Lee et al. who demonstrated the improved sensitivity to a nephrotoxicant in kidney organoids cultured under flow.⁶⁶ However, to date, the ability to address and perfuse the organoid microvasculature from larger vessels has yet to be demonstrated.

Creating kidney organoids with a perfusable vasculature *in vivo* can be achieved by implantation into mice;^{8,4,9,69,85,98,112} or chick chorioallantoic membranes,³⁷ where host vascular integration ensues leading to both organoid maturation and/or rudimentary filtration. Organ-on-chip methods may allow organoids to mimic these native processes *in vitro* through vasculature recruitment from a nearby endothelialized channel or integration with prepatterned vasculature.¹¹⁶ For example, Nashimoto et al. demonstrated that placing tumor spheroids between endothelialized channels induced vasculature sprouting and fusing leading to perfusable microvasculature between the tumor spheroids and channels.⁷⁷ While Zhang et al. 3D printed a vascular scaffold that was subsequently seeded with cardiomyocytes to produce vascularized cardiac tissue.¹¹⁷ While these are encouraging *in vitro* demonstrations, neither of these approaches have been extended to kidney organoids. Moreover, we posit that vasculature recruitment may offer a better approach, since it mimics *in vivo* vascularization of kidney organoids.¹¹⁶

Once perfusable, vascularized kidney organoids are obtained *in vitro*, the next crucial step will be the introduction and integration of a collecting duct system that allows for primitive filtrate removal. Current differentiation protocols that yield MM-derived cell populations lack UB-derived collecting ducts.^{22,75,93} However, when embryonic MM and UB populations are combined *in vitro* via core-shell assembly, one observes the formation of a hierarchical collecting duct system.^{36,101} As an example, Tanigawa and Tanaka et al. have demonstrated this for murine iPSC-derived kidney organoids by combining induced MM, UB and stromal populations.⁹⁶ The assembly of murine ESC-derived NPs, UBs, and stromal progenitors resulted in the generation of organ-like kidney structure, which comprise nephrons with multiple types of stromal cells accumulated around branching UBs.⁹⁶ However, to our knowledge, this has yet to be achieved in human kidney organoids.

The ability to achieve filtrate drainage in kidney organoids *in vitro* is important for two reasons: (1) absent this capability, filtrate accumulation will lead to hydronephrosis in vascularized organoids,¹¹⁵ and (2) filtrate collection is essential for functional assessment. At present, fluorescent probes can be used to study filtration and transporter processes,^{33,8,59} while micro-puncture techniques enable filtrate composition to be assessed in individual tubules.^{15,103} To generate kidney organoids with an open collecting duct or ureter, one can draw inspiration from kidney development and *in vitro* vasculature recruitment models. We hypothesize that such organoids might be generated by embedding them within an extracellular matrix *in vitro* that

contains an open nephric duct, produced for example by bioprinting, from which a UB arises and invades a surrounding MM.

Engineering kidney tissues. By further advancing kidney organoids into functional models, one can unlock their potential as building blocks for engineering kidney tissue for therapeutic use. Several 3D bioprinting methods enable the fabrication of human tissues using organoids or multicellular spheroids as building blocks.^{3,26,41,65,89} For example, Lawlor and Vanslam-brouck et al. created two-dimensional kidney tissue sheets composed of closely printed lines of IM cells that fuse together and further develop over time. The resulting kidney tissue demonstrated correctly segmented nephrons, including proximal tubules that exhibited albumin uptake.⁶⁵ Another promising method to generating vascularized human tissues, known as sacrificial writing in functional tissue (SWIFT), was recently developed by Skylar-Scott and Uzel et al.⁸⁹ This method involves printing branched vascular networks within a living tissue matrix comprised of densely cellular, organoid building blocks. By embedding a perfusable vasculature within these tissues, one can overcome the nutrient diffusion limitation that gives rise to core necrosis and fibrosis in organoids with diameters above $\sim 500 \mu\text{m}$, resulting in a large viable tissue.^{89,116} Since SWIFT biomanufacturing is broadly applicable, this method would enable the construction of vascularized kidney tissues using organoids as organ building blocks. In the future, the large scale production of the vascularized kidney tissues from hundreds of thousands of organoids within these engineered tissues will ultimately enable the generation of engineered organ-like structures of kidney containing 1 M+ nephrons akin to human kidneys.^{11,78}

Kidney organoids to overcome post-transplant immune rejection. Kidney transplantation is currently the optimal treatment for ESKD patients; however, rejection of the allogeneic kidney transplants by the host immune system is a major obstacle, leading to graft loss.^{50,83} Rejection is caused by an immune response against the transplanted organ, driven by the recognition of the transplanted kidney tissue antigens as foreign bodies by the host immune system. Immune rejection of the transplanted graft is classified as hyperacute, acute, and chronic rejection, based on the histopathological features and the time frame of the rejection process following the transplantation.¹ In the course of xenotransplantation from pigs or nonhuman primates to humans, both hyperacute and acute humoral or acute cellular rejection would be evoked by the circulating antibodies, complement, neutrophils, natural killer (NK) cells, T cells, and macrophages.⁷⁰

Hyperacute rejection (HAR) is characterized by extensive thrombotic microangiopathy (TMA),^{1,70} mediated by multiple factors including pre-existing IgM/IgG alloantibodies of the recipient to the porcine carbohydrate epitopes (Gal α 1–3Gal β 1–4GlcNAc: α 1,3Gal epitope), complement activation, and the hypercoagulable state.^{35,114} Recently, 2 studies attempted to overcome the hyperacute rejection by knocking-out α -1,3-galactosyltransferase enzyme and modifying multiple genes associated with complement, coagulant, and immune regulation in the setting of porcine kidney xenotransplantation to brain-dead human recipients.^{72,80} However, it is still unclear if those gene editing can prevent TMA in humans for the long-term. In contrast, this hyperacute rejection does not constitute an obstacle for the possible transplantation of human iPSC-derived kidney organoids as human kidney tissues do not express α 1,3Gal epitope.

Acute and chronic rejection involve damage to both the graft parenchyma and blood vessels, mediated by alloreactive T cells and alloantibodies against the human leukocyte antigens (HLAs) expressed by the donor organs and/or donor APCs.^{1,52,104} HLAs are the predominant target of the immune system based on its great heterogeneity and polymorphism between individuals.⁷³ The degree of HLA mismatch between donor and recipient correlates with the graft survival as the presence of higher mismatches is associated with a significant higher risk of graft failure.¹⁰⁷ With the future advancement of 3D-biofabricated kidney structures derived from HLA-edited hiPSCs via CRISPR/Cas9 technology, it is conceivable to generate immune-tolerant kidney tissues with manipulated HLA molecules, which enhance the compatibility between donors and recipients, thereby reducing the risk of acute and/or chronic rejection responses. This novel approach may also dampen the need and dosage of current immunosuppressive agents such as calcineurin inhibitors (cyclosporine, tacrolimus), mammalian target of rapamycin (*mTOR*) inhibitors, and T cell-depleting induction therapy.^{23,79} Furthermore, enhancing the expression of coinhibitory receptors on kidney cells may provide an additional protection against the immune response to suppress rejection,⁸² which might translate into the ultimate personalized therapeutic strategy in transplantation.

Recently, Cowan et al. generated hypoimmunogenic hPSCs using the CRISPR/Cas9 system by ablation of HLA-A/B/C and HLA class II, and overexpression of inhibitory signals such as PD-L1, HLA-G, and CD47 to suppress T cell, NK cell responses, and macrophage engulfment.⁵⁰ However, overexpression of checkpoint molecules like PD-L1 bears the risk of malignancy by promoting tumor escape from the immune surveillance.⁵¹ In another study, Hotta et al. developed immune compatible iPSCs by preserving HLA-C,

inhibitory for NK cells, in the absence of HLA-A/B and CIITA [major histocompatibility complex (MHC) class II transactivator essential for the transcription of class II HLA genes].¹¹³ Consequently, 12 HLA-edited (HLA-A- and -B-deleted and HLA-C-retained) iPSC lines could serve as immunologically compatible iPSCs for nearly >90% of the world population,¹¹³ thereby enabling the generation of universal kidney organoids as building blocks, which can give rise to immune-compatible organotypic kidney structures in the future. Overall, the development of these broadly compatible kidney structures may allow a larger scale and cost-effective therapeutic strategy as an alternative to the generation of patient-specific kidney organoids.

Current limitations of kidney organoid models. Kidney organoids offer a tremendous opportunity for the future medical applications, such as generation of immune compatible bioengineered kidneys for the growing demand from CKD patients. However, this unique technology comes with its own restrictions. When a previous study compared the maturation status of organoids directly differentiated via the Morizane and Takasato protocols by using the single-cell transcriptomics of fetal and adult kidney cells, they demonstrated that the organoids by both protocols gave rise to mostly immature cell types at early differentiation stages (~26 days) with respect to their adult counterparts.¹¹⁰ Even though the epithelial cell types of the adult kidney and kidney organoids correlated well, organoid-derived cells tended to express developmental genes. Efforts need to be made to extend the organoid culture for further maturation. In addition, the single-cell transcriptomics suggested the protocol differences. Morizane protocol appears to induce more matured nephron epithelia than Takasato protocol according to Wu's study. Although it is difficult to have a definitive conclusion due to batch-to-batch variations, further improvements on the current protocols should also be considered. Furthermore, it is essential to improve the current methodology for the evaluation of functional assessment of kidney organoids, which are building blocks for the in vitro bioengineered kidneys.⁸⁴ Therefore, the organoids from established protocols should be tested to confirm the recapitulation of normal kidney functions such as glomerular filtration, tubular reabsorption/secretion, and production of erythropoietin and angiotensin II hormones.

CONCLUSION

The kidney functions as a critical organ maintaining the homeostasis of body fluid and electrolyte status by effectively removing the metabolic waste products. Acute kidney injury or progressive chronic kidney

disease may result in renal insufficiency with serious systemic complications in the patients, who eventually need to undergo renal replacement therapies such as dialysis and transplantation, thereby leading to a high socioeconomical burden in the healthcare system. Conventional research techniques remain insufficient to fully recapitulate the kidney physiology and pathogenesis underlying the complex kidney diseases. Thus, the development of directed differentiation protocols, which simulate *in vivo* nephrogenesis, enabled the generation of *in vitro* 3D kidney organoids from hPSCs. Even though the application of patient iPSC-derived kidney organoids advanced our understanding of the kidney maturation, injury, and repair, we still need to facilitate the maturation of glomerular vascularization and incorporate collecting duct system into organoids by modifying the current organoid protocol with additional novel techniques such as 3D-bioprinting and organ-on-a-chip. The ultimate technology of organoids-on-chip may provide the better translational utility for disease modeling and drug screening, potentially replacing the animal models in the future.

ACKNOWLEDGMENTS

Conflicts of Interest: All authors have read the journal's policy on disclosure of potential conflicts of interest. J.A.L. and R.M. are co-inventors on patents (PCT/US2018/036677) that are assigned to President and Fellows of Harvard College and Mass General Brigham. J.A.L. and R.M. serve on the Scientific Advisory Board of Trestle Biotherapeutics, which has licensed technology from their labs. J.A.L. is a cofounder of AcousticaBio and serves as a scientific advisor for Autodesk, Azul3D, Desktop Health (a subsidiary of Desktop Metal), Mooji Meats, and Trestle Biotherapeutics. S.G.M.U., and J.A.L. are inventors of the SWIFT biomanufacturing method and have filed a patent on this work.

The authors thank the following funding that supported this work: NIH DP2EB029388 award (R.M.), NIH U01EB028899/DK127587 grant (J.A.L., R.M.), NIH UC2DK126023 grant (J.A.L., R.M.), NIH UH3TR002155 grant (J.A.L., R.M.), NWO 019.201EN.005 grant by Dutch Research Council (R.C.G.), Wellcome Leap HOPE Program (L.V.R., J.A.L., R.M.).

All authors have read the journal's authorship agreement and that the manuscript has been reviewed by and approved by all named authors. Authors thank the lab members of Riella, Lewis and Morizane labs for insightful discussion, Kenichi Kobayashi for his supervision during kidney organoid generation, and Annie Moisan, HOPE Program Director at Wellcome Leap, for interactive discussion about the generation of *in vitro* 3D kidney tissues derived from immune compatible organoid building blocks.

REFERENCES

1. Abbas, A. K., Andrew, Lichtman, & Pillai, Shiv. H. (2012). Cellular and molecular immunology (7th ed.). (Philadelphia, PA: Elsevier Saunders).
2. Andersen Monica L, Winter Lucile MF. Animal models in biological and biomedical research – experimental and ethical concerns. *An Acad Bras Cienc* 2019;91(1):1–14. <https://doi.org/10.1590/0001-3765201720170238.1>.
3. Ayan Bugra, Heo Dong Nyoung, Zhang Zhifeng, Dey Madhuri, Povilianskas Adomas, Drapaca Corina, et al. Aspiration-assisted bioprinting for precise positioning of biologics. *Sci Adv* 2020;6(10):1–17. <https://doi.org/10.1126/sciadv.aaw5111>.
4. Bantounas Ioannis, Ranjzad Parisa, Tengku Faris, Silajdzic Edina, Forster Duncan, Asselin Marie Claude, et al. Generation of functioning nephrons by implanting human pluripotent stem cell-derived kidney progenitors. *Stem Cell Rep* 2018;10(3):766–79. <https://doi.org/10.1016/j.stemcr.2018.01.008>.
5. Bar-Ephraim Yotam E, Kretzschmar Kai, Clevers Hans. Organoids in immunological research. *Nature Rev Immunol* 2020;20(5):279–93. <https://doi.org/10.1038/s41577-019-0248-y>.
6. Barak Hila, Huh Sung Ho, Chen Shuang, Jeanpierre Cécile, Martinovic Jelena, Parisot Mélanie, et al. FGF9 and FGF20 maintain the stemness of nephron progenitors in mice and man. *Develop Cell* 2012;22(6):1191–207. <https://doi.org/10.1016/j.devcel.2012.04.018>.
7. Bennett Christina N, Ross Sarah E, Longo Kenneth A, Bajnok Laszlo, Hemati Nahid, Johnson Kirk W, et al. Regulation of Wnt Signaling during Adipogenesis. *J Biol Chem* 2002;277(34):30998–4. <https://doi.org/10.1074/jbc.M204527200>.
8. Berg Cathelijne WVan Den, Koudijs Angela, Ritsma Laila, Rabelink Ton J. In vivo assessment of size-selective glomerular sieving in transplanted human induced pluripotent stem cell-derived kidney organoids. *J Am Society of Nephrol* 2020;31(5):921–9. <https://doi.org/10.1681/ASN.2019060573>.
9. Berg Cathelijne Wvan den, Ritsma Laila, Cristina Avramut M, Wiersma Loes E, Berg Bernard Mvan den, Leuning Daniëlle G, et al. Renal subcapsular transplantation of PSC-derived kidney organoids induces neo-vasculogenesis and significant glomerular and tubular maturation *in vivo*. *Stem Cell Rep* 2018;10(3):751–65. <https://doi.org/10.1016/j.stemcr.2018.01.041>.
10. Bergmann Carsten, Guay-Woodford Lisa M, Harris Peter C, Horie Shigeo, Peters Dorian JM, Torres Vicente E. Polycystic kidney disease. *Nature Rev Dis Primers* 2018;4(1). <https://doi.org/10.1038/s41572-018-0047-y>.
11. Bertram John F, Douglas-Denton Rebecca N, Diouf Boucar, Hughson Michael D, Hoy Wendy E. Human nephron number: implications for health and disease. *Pediatric Nephrol* 2011;26(9):1529–33. <https://doi.org/10.1007/s00467-011-1843-8>.
12. Beura Lalit K, Hamilton Sara E, Bi Kevin, Schenkel Jason M, Odumade Oludare A, Casey Kerry A, et al. Normalizing the environment recapitulates adult human immune traits in laboratory mice. *Nature* 2016;532(7600):512–6. <https://doi.org/10.1038/nature17655>.
13. Braun, Daniela A., & Hildebrandt, Friedhelm (2022). Ciliopathies.
14. Breda Philippe Christophe, Wiech Thorsten, Meyer-Schwesinger Catherine, Grahmmer Florian, Huber Tobias, Panzer

- Ulf, et al. Renal proximal tubular epithelial cells exert immunomodulatory function by driving inflammatory CD4+ T cell responses. *Am J Physiol - Renal Physiol* 2019;317(1):F77–89. <https://doi.org/10.1152/ajprenal.00427.2018>.
15. Burg M, Grantham J, Abramow M, Orloff J. Preparation and study of fragments of single rabbit nephrons. *Am J Physiol* 1966;210(6):1293–8. <https://doi.org/10.1152/ajplegacy.1966.210.6.1293>.
 16. Carroll Thomas J, Das Amrita. Defining the signals that constitute the nephron progenitor niche. *J Am Society of Nephrol* 2013;24(6):873–6. <https://doi.org/10.1681/ASN.2012090931>.
 17. Carroll Thomas J, Park Joo Seop, Hayashi Shigemi, Majumdar Arindam, McMahon Andrew P. Wnt9b plays a central role in the regulation of mesenchymal to epithelial transitions underlying organogenesis of the mammalian urogenital system. *Develop Cell* 2005;9(2):283–92. <https://doi.org/10.1016/j.devcel.2005.05.016>.
 18. Choudhury Mohammad Ikbal, Li Yizeng, Mistriotis Panagiotis, Vasconcelos Ana Carina N, Dixon Eryn E, Yang Jing, et al. Kidney epithelial cells are active mechano-biological fluid pumps. *Nature Commun* 2022;13(1):1–13. <https://doi.org/10.1038/s41467-022-29988-w>.
 19. Chun, Justin, Riella, Cristian V., Chung, Hyunjae, Shah, Shrijal S., Wang, Minxian, Magraner, Jose M., et al. (2022). DGAT2 inhibition potentiates lipid droplet formation to reduce cytotoxicity in APOL1 kidney risk variants. *J Am Society of Nephrol : JASN* 33 (5), 889–907. doi: 10.1681/ASN.2021050723
 20. Clevers Hans. Modeling development and disease with organoids. *Cell* 2016;165(7):1586–97. <https://doi.org/10.1016/j.cell.2016.05.082>.
 21. Coelho Silvia, Cabral Guadalupe, Lopes José A, Jacinto António. Renal regeneration after acute kidney injury. *Nephrology* 2018;23(9):805–14. <https://doi.org/10.1111/nep.13256>.
 22. Combes Alexander N, Phipson Belinda, Lawlor Kynan T, Dorison Aude, Patrick Ralph, Zappia Luke, et al. Correction: single cell analysis of the developing mouse kidney provides deeper insight into marker gene expression and ligand-receptor cross-talk. *Develop (Cambridge)* 2019;(12):146. <https://doi.org/10.1242/dev.182162>.
 23. Cooper James E. Evaluation and treatment of acute rejection in kidney allografts. *Clin J Am Society of Nephrol* 2020;15(3):430–8. <https://doi.org/10.2215/CJN.11991019>.
 24. Costantini Frank, Kopan Raphael. Patterning a complex organ: branching morphogenesis and nephron segmentation in kidney development. *Develop Cell* 2010;18(5):698–712. <https://doi.org/10.1016/j.devcel.2010.04.008>.
 25. Cruz Nelly M, Song Xuwen, Czerniecki Stefan M, Gulieva Ramila E, Churchill Angela J, Kim Yong Kyun, et al. Organoid cystogenesis reveals a critical role of microenvironment in human polycystic kidney disease. *Nature Mater* 2017;16(11):1112–9. <https://doi.org/10.1038/NMAT4994>.
 26. Daly Andrew C, Davidson Matthew D, Burdick Jason A. 3D bioprinting of high cell-density heterogeneous tissue models through spheroid fusion within self-healing hydrogels. *Nature Commun* 2021;12(1):1–13. <https://doi.org/10.1038/s41467-021-21029-2>.
 27. Deschamps Jacqueline, Nes Johan van. Developmental regulation of the hox genes during axial morphogenesis in the mouse. *Development* 2005;132(13):2931–42. <https://doi.org/10.1242/dev.01897>.
 28. Dong Lihua, Pietsch Stefan, Englert Christoph. Towards an understanding of kidney diseases associated with WT1 mutations. *Kidney Internatl* 2015;88(4):684–90. <https://doi.org/10.1038/ki.2015.198>.
 29. Dressler, Gregory R. (2009). Advances in early kidney specification, development and patterning. *Development* 136 (23), 3863–74. doi: 10.1242/dev.034876
 30. Duval, Kayla, Grover, Hannah, Han, Li Hsin, Mou, Yongchao, Pegoraro, Adrian F., Fredberg, Jeffery, et al. (2017). Modeling physiological events in 2D vs. 3D Cell culture. *Physiology* 32 (4), 266–77. doi: 10.1152/physiol.00036.2016
 31. Ettou Sandrine, Jung Youngsook L, Miyoshi Tomoya, Jain Dhawal, Hiratsuka Ken, Schumacher Valerie, et al. Epigenetic transcriptional reprogramming by WT1 mediates a repair response during podocyte injury. *Sci Adv* 2020;6(30):1–14. <https://doi.org/10.1126/sciadv.abb5460>.
 32. Forbes Thomas A, Howden Sara E, Lawlor Kynan, Phipson Belinda, Maksimovic Jovana, Hale Lorna, et al. Patient-IPSC-derived kidney organoids show functional validation of a ciliopathic renal phenotype and reveal underlying pathogenetic mechanisms. *Am J Human Genetics* 2018;102(5):816–31. <https://doi.org/10.1016/j.ajhg.2018.03.014>.
 33. Freedman Benjamin S, Brooks Craig R, Lam Albert Q, Fu Hongxia, Morizane Ryuji, Agrawal Vishesh, et al. Modelling kidney disease with CRISPR-mutant kidney organoids derived from human pluripotent epiblast spheroids. *Nature Commun* 2015;6. <https://doi.org/10.1038/ncomms9715>.
 34. Freire Analía V, Ropelato María Gabriela, Rey Rodolfo A. Development and function of the ovaries and testes in the fetus and neonate. *Maternal-Fetal and Neonatal Endocrinol: Physiol, Pathophysiol, and Clin Managt* 2019;625–41. <https://doi.org/10.1016/B978-0-12-814823-5.00037-4>.
 35. Ganchiku, Yoshikazu, & Riella, Leonardo V. (2022). Pig-to-human kidney transplantation using brain-dead donors as recipients: one giant leap, or only one small step for transplantkind? *Xenotransplantation*, no. April (May): 1–6. doi: 10.1111/xen.12748
 36. Ganeva Veronika, Unbekandt Mathieu, Davies Jamie A. An improved kidney dissociation and re-aggregation culture system results in nephrons arranged organotypically around a single collecting duct system. *Organogenesis* 2011;7(2):83–7. <https://doi.org/10.4161/org.7.2.14881>.
 37. Garreta Elena, Prado Patricia, Tarantino Carolina, Oria Roger, Fanlo Lucia, Martí Elisa, et al. Fine tuning the extracellular environment accelerates the derivation of kidney organoids from human pluripotent stem cells. *Nature Materials* 2019;18(4):397–405. <https://doi.org/10.1038/s41563-019-0287-6>.
 38. Gaunt Stephen J, George Martin, Paul Yu Lee. Direct activation of a mouse Hoxd11 axial expression enhancer by Gdf11/SMAD signaling. *Develop Biol* 2013;383(1):52–60. <https://doi.org/10.1016/j.ydbio.2013.08.025>.
 39. Genovese Giulio, Friedman David J, Ross Michael D, Lecordier Laurence, Uzureau Pierrick, Freedman Barry I, et al. Association of trypanolytic ApoL1 variants with kidney disease in African Americans. *Science* 2010;329(5993):841–5. <https://doi.org/10.1126/science.1193032>.
 40. Goldwich Andreas, Burkard Miriam, Ölke Martha, Daniel Christoph, Amann Kerstin, Hugo Christian, et al. Podocytes are nonhematopoietic professional antigen-presenting cells. *J Am Society of Nephrol* 2013;24(6):906–16. <https://doi.org/10.1681/ASN.2012020133>.
 41. Goulart Ernesto, Caires-Junior Luiz Carlos de, Telles-Silva Kayque Alves, Araujo Bruno Henrique Silva, Rocco Silvana Aparecida, Sforca Mauricio, et al. 3D bioprinting of liver spheroids derived from human induced pluripotent stem cells

- sustain liver function and viability in vitro. *Biofabrication* 2020;12:015010.
42. Grahammer Florian, Schell Christoph, Huber Tobias B. The podocyte slit diaphragm—from a thin grey line to a complex signaling hub. *Nature Rev Nephrol* 2013;9(10):587–98. <https://doi.org/10.1038/nrneph.2013.169>.
 43. Gupta Navin, Dilmen Emre, Morizane Ryuji. 3D kidney organoids for bench-to-bedside translation. *J Molecular Med* 2021;99(4):477–87. <https://doi.org/10.1007/s00109-020-01983-y>.
 44. Gupta Navin, Matsumoto Takuya, Hiratsuka Ken, Garcia Saiz Edgar, Galichon Pierre, Miyoshi Tomoya, et al. Modeling injury and repair in kidney organoids reveals that homologous recombination governs tubular intrinsic repair. *Sci Transl Med* 2022;14(634):1–16. <https://doi.org/10.1126/scitranslmed.abj4772>.
 45. Gupta, Navin, & Morizane, Ryuji (2021). Kidney development to kidney organoids and back again. *Seminars in Cell and Developmental Biology*, no. September. doi: 10.1016/j.semedb.2021.09.017
 46. Gupta Navin, Susa Koichiro, Morizane Ryuji. Regenerative medicine, disease modeling, and drug discovery in human pluripotent stem cell-derived kidney tissue. *Euro Med J. Reproductive Health* 2017;3(1):57–67, Available at: <http://www.ncbi.nlm.nih.gov/pubmed/31157117> <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC6544146>.
 47. Hakrroush Samy, Kopp Sarah Birgit, Tampe Désirée, Gersmann Ann Kathrin, Korsten Peter, Zeisberg Michael, et al. Variable expression of programmed cell death protein 1-ligand 1 in kidneys independent of immune checkpoint inhibition. *Frontiers in Immunol* 2021;11:1–7. <https://doi.org/10.3389/fimmu.2020.624547> (January)doi:.
 48. Hale Lorna J, Howden Sara E, Phipson Belinda, Lonsdale Andrew, Er Pei X, Ghobrial Irene, et al. 3D organoid-derived human glomeruli for personalised podocyte disease modelling and drug screening. *Nature Commun* 2018;9(1). <https://doi.org/10.1038/s41467-018-07594-z>.
 49. Hamilton Sara E, Badovinac Vladimir P, Beura Lalit K, Pierson Mark, Jameson Stephen C, Masopust David, et al. New insights into the immune system using dirty mice. *The J Immunol* 2020;205(1):3–11. <https://doi.org/10.4049/jimmunol.2000171>.
 50. Han Xiao, Wang Mengning, Duan Songwei, Franco Paul J, Kenty Jennifer Hyeje Ryu, Hedrick Preston, et al. Generation of hypoinmunogenic human pluripotent stem cells. *Proceedings of the National Academy of Sciences of the United States of America* 2019;116(21):10441–6. <https://doi.org/10.1073/pnas.1902566116>.
 51. He Jiabei, Hu Ying, Hu Mingming, Li Baolan. Development of PD-1/PD-L1 pathway in tumor immune microenvironment and treatment for non-small cell lung cancer. *Scientific Reports* 2015;5:1–9. <https://doi.org/10.1038/srep13110>.
 52. Heemann Uwe, Lutz Jens. Pathophysiology and treatment options of chronic renal allograft damage. *Nephrol Dialysis Transplantation* 2013;28(10):2438–46. <https://doi.org/10.1093/ndt/gft087>.
 53. Hill Nathan R, Fatoba Samuel T, Oke Jason L, Hirst Jennifer A, O'Callaghan Christopher A, Lasserson Daniel S, et al. Global prevalence of chronic kidney disease – a systematic review and meta-analysis Edited by Giuseppe Remuzzi *PLOS ONE* 2016;11(7):1–18. <https://doi.org/10.1371/journal.pone.0158765>.
 54. Hoefele Julia, Mayer Karin, Scholz Manuela, Klein Hanns Georg. Novel PKD1 and PKD2 mutations in Autosomal Dominant Polycystic Kidney Disease (ADPKD). *Nephrol Dialysis Transplantation* 2011;26(7):2181–8. <https://doi.org/10.1093/ndt/gfq720>.
 55. Homan Kimberly A, Gupta Navin, Kroll Katharina T, Kolesky David B, Skylar-Scott Mark, Miyoshi Tomoya, et al. Flow-enhanced vascularization and maturation of kidney organoids in vitro. *Nature Methods* 2019;16(3):255–62. <https://doi.org/10.1038/s41592-019-0325-y>.
 56. Huber Tobias B, Benzing Thomas. The Slit diaphragm: a signaling platform to regulate podocyte function. *Curr Opin Nephrol and Hypertension* 2005;14(3):211–6. <https://doi.org/10.1097/01.mnh.0000165885.85803.a8>.
 57. Humphreys Benjamin D, Todd Valerius M, Kobayashi Akio, Mugford Joshua W, Soeung Savuth, Duffield Jeremy S, et al. Intrinsic epithelial cells repair the kidney after injury. *Cell Stem Cell* 2008;2(3):284–91. <https://doi.org/10.1016/j.stem.2008.01.014>.
 58. Imura Tadahiro, Pourquié Olivier. Collinear activation of Hoxb genes during gastrulation is linked to mesoderm cell ingression. *Nature* 2006;442(7102):568–71. <https://doi.org/10.1038/nature04838>.
 59. Jansen J, Fedecostante M, Wilmer MJ, Peters JG, Kreuser UM, van den Broek PH, et al. Bioengineered kidney tubules efficiently excrete uremic toxins. *Scientific Reports* 2016;6(1):26715. <https://doi.org/10.1038/srep26715>.
 60. Jourde-Chiche Noémie, Fakhouri Fadi, Dou Laetitia, Bellien Jeremy, Burtay Stéphane, Frimat Marie, et al. Endothelium structure and function in kidney health and disease. *Nature Reviews Nephrol* 2019;15(2):87–108. <https://doi.org/10.1038/s41581-018-0098-z>.
 61. Seok Junhee, Warren HShaw, Cuenca Alex G, Mindrinos Michael N, Baker Henry V, Xu Weihong, et al. Genomic responses in mouse models poorly mimic human inflammatory diseases. *Proceedings of the National Academy of Sciences of the United States of America* 2013;110(9):3507–12. <https://doi.org/10.1073/pnas.1222878110>.
 62. Kaku Yusuke, Taguchi Atsuhiko, Tanigawa Shunsuke, Haque Fahim, Sakuma Tetsushi, Yamamoto Takashi, et al. PAX2 is dispensable for in vitro nephron formation from human induced pluripotent stem cells. *Scientific Reports* 2017;7(1):1–12. <https://doi.org/10.1038/s41598-017-04813-3>.
 63. Karner Courtney M, Das Amrita, Ma Zhendong, Self Michelle, Chen Chuo, Lum Lawrence, et al. Canonical Wnt9b signaling balances progenitor cell expansion and differentiation during kidney development. *Development* 2011;138(7):1247–57. <https://doi.org/10.1242/dev.057646>.
 64. Kuppe Christoph, Ibrahim Mahmoud M, Kranz Jennifer, Zhang Xiaoting, Ziegler Susanne, Perales-Patón Javier, et al. Decoding myofibroblast origins in human kidney fibrosis. *Nature* 2021;589(7841):281–6. <https://doi.org/10.1038/s41586-020-2941-1>.
 65. Lawlor Kynan T, Vanslambrouck Jessica M, William Higgins J, Chambon Alison, Bishard Kristina, Arndt Derek, et al. Cellular extrusion bioprinting improves kidney organoid reproducibility and conformation. *Nature Materials* 2021;20(2):260–71. <https://doi.org/10.1038/s41563-020-00853-9>.
 66. Lee Han Na, Choi Yoon Young, Kim Jin Won, Lee Young Seo, Choi Ji Wook, Kang Taewook, et al. Effect of biochemical and biomechanical factors on vascularization of kidney organoid-on-a-chip. *Nano Convergence* 2021;8(1). <https://doi.org/10.1186/s40580-021-00285-4>.
 67. Lemos Dario R, McMurdo Michael, Karaca Gamze, Wilflingseder Julia, Leaf Irina A, Gupta Navin, et al. Interleukin-1b activates a MYC-dependent metabolic switch in kidney stromal

- cells necessary for progressive tubulointerstitial fibrosis. *J Am Society of Nephrol* 2018;29(6):1690–705. <https://doi.org/10.1681/ASN.2017121283>.
68. Little Melissa H, Combes Alexander N. Kidney organoids: accurate models or fortunate accidents. *Genes and Development* 2019;33(19–20):1319–45. <https://doi.org/10.1101/gad.329573.119>.
 69. Low Jian Hui, Li Pin, Chew Elaine Guo Yan, Zhou Bingrui, Suzuki Keiichiro, Zhang Tian, et al. Generation of human PSC-derived kidney organoids with patterned nephron segments and a de novo vascular network. *Cell Stem Cell* 2019;25(3):373–87. <https://doi.org/10.1016/j.stem.2019.06.009>.
 70. Lu Tianyu, Yang Bochao, Wang Ruolin, Qin Chuan. Xenotransplantation: current status in preclinical research. *Frontiers in Immunol* 2020;10. <https://doi.org/10.3389/fimmu.2019.03060> January.
 71. Maeshima Akito, Yamashita Shin, Maeshima Kyoko, Kojima Itaru, Nojima Yoshihisa. Activin a produced by ureteric bud is a differentiation factor for metanephric mesenchyme. *J Am Society of Nephrol* 2003;14(6):1523–34. <https://doi.org/10.1097/01.ASN.0000067419.86611.21>.
 72. Montgomery Robert A, Stern Jeffrey M, Lonze Bonnie E, Tata-pudi Vasishtha S, Mangiola Massimo, Wu Ming, et al. Results of two cases of pig-to-human kidney xenotransplantation. *New Eng J Med* 2022;386(20):1889–98. <https://doi.org/10.1056/nejmoa2120238>.
 73. Montgomery Robert A, Tatapudi Vasishtha S, Leffell Mary S, Zachary Andrea A. HLA in transplantation. *Nature Reviews Nephrol* 2018;14(9):558–70. <https://doi.org/10.1038/s41581-018-0039-x>.
 74. Morizane Ryuji, Bonventre Joseph V. Generation of nephron progenitor cells and kidney organoids from human pluripotent stem cells. *Nature Protocols* 2017;12(1):195–207. <https://doi.org/10.1038/nprot.2016.170>.
 75. Morizane Ryuji, Lam Albert Q, Freedman Benjamin S, Kishi Seiji, Todd Valerius M, Bonventre Joseph V. Nephron organoids derived from human pluripotent stem cells model kidney development and injury. *Nature Biotechnol* 2015;33(11):1193–200. <https://doi.org/10.1038/nbt.3392>.
 76. Mugford Joshua W, Sipilä Petra, McMahon Jill A, McMahon Andrew P. Oslr expression demarcates a multi-potent population of intermediate mesoderm that undergoes progressive restriction to an Oslr-dependent nephron progenitor compartment within the mammalian kidney. *Developmental Biol* 2008;324(1):88–98. <https://doi.org/10.1016/j.ydbio.2008.09.010>.
 77. Nashimoto Yuji, Okada Ryu, Hanada Sanhiro, Arima Yui-chiro, Nishiyama Koichi, Miura Takashi, et al. Vascularized cancer on a chip: the effect of perfusion on growth and drug delivery of tumor spheroid. *Biomaterials* 2020;229:119547. <https://doi.org/10.1016/j.biomaterials.2019.119547> October 2019.
 78. Nishinakamura Ryuichi. Human kidney organoids: progress and remaining challenges. *Nature Reviews Nephrol* 2019;15(10):613–24. <https://doi.org/10.1038/s41581-019-0176-x>.
 79. Pascual Julio, Berger Stefan P, Witzke Oliver, Tedesco Helio, Mulgaonkar Shamkant, Qazi Yasir, et al. Everolimus with reduced calcineurin inhibitor exposure in renal transplantation. *J Am Society of Nephrol* 2018;29(7):1979–91. <https://doi.org/10.1681/asn.2018010009>.
 80. Porrett Paige M, Orandi Babak J, Kumar Vineeta, Houpp Julie, Anderson Douglas, Cozette Killian A, et al. First clinical-grade porcine kidney xenotransplant using a human decedent model. *Am J Transplantation* 2022;22(4):1037–53. <https://doi.org/10.1111/ajt.16930>.
 81. Potter SSteven, Hartman Heather A, Kwan Kin M, Behringer Richard R, Patterson Larry T. Laser capture-microarray analysis of *lim1* mutant kidney development. *Genesis* 2007;45(7):432–9. <https://doi.org/10.1002/dvg.20309>.
 82. Riella LV, Paterson AM, Sharpe AH, Chandraker A. Role of the PD-1 pathway in the immune response. *Am J Transplantation* 2012;12(10):2575–87. <https://doi.org/10.1111/j.1600-6143.2012.04224.x>.
 83. Riella Leonardo V, Djamali Arjang, Pascual Julio. Chronic allograft injury: mechanisms and potential treatment targets. *Transplantation Rev* 2017;31(1):1–9. <https://doi.org/10.1016/j.trre.2016.10.005>.
 84. Rizki-Safitri, Astia, Traitteur, Tamara, & Morizane, Ryuji (2021). Bioengineered kidney models: methods and functional assessments. *Function* 2(4): 1–17. doi: 10.1093/function/zqab026
 85. Ryan Anne R, England Alicia R, Chaney Christopher P, Cowdin Mitzy A, Hiltabidle Max, Daniel Edward, et al. Vascular deficiencies in renal organoids and ex vivo kidney organogenesis. *Developmental Biol* 2021;477:98–116. <https://doi.org/10.1016/j.ydbio.2021.04.009> Aprildoi:.
 86. Saxen, Lauri. (1987). *Organogenesis of the Kidney*. Cambridge University Press. doi: 10.1017/CBO9780511565083
 87. Schell C, Wanner N, Huber TB. Glomerular development - shaping the multi-cellular filtration unit. *Seminars in Cell and Developmental Biol* 2014;36:39–49. <https://doi.org/10.1016/j.semcdb.2014.07.016>.
 88. Sharp Cierra N, Doll Mark A, Megyesi Judit, Oropilla Gabrielle B, Beverly Levi J, Siskind Leah J. Subclinical kidney injury induced by repeated cisplatin administration results in progressive chronic kidney disease. *Am J Physiol - Renal Physiol* 2018;315(1):F161–72. <https://doi.org/10.1152/ajprenal.00636.2017>.
 89. Skylar-Scott Mark A, Uzel Sebastien GM, Nam Lucy L, Ahrens John H, Truby Ryan L, Damaraju Sarita, et al. Biomaterial manufacturing of organ-specific tissues with high cellular density and embedded vascular channels. *Sci Advances* 2019;5(9):1–13. <https://doi.org/10.1126/sciadv.aaw2459>.
 90. Solanki Ashish K, Arif Ehtesham, Srivastava Pankaj, Furcht Christopher M, Rahman Bushra, Wen Pei, et al. Phosphorylation of slit diaphragm proteins NEPHRIN and NEPH1 upon binding of HGF promotes podocyte repair. *J Biol Chemistry* 2021;297(3):101079. <https://doi.org/10.1016/j.jbc.2021.101079>.
 91. Starke Astrid, Lindenmeyer Maja T, Segerer Stephan, Neusser Matthias A, Rüsi Barbara, Schmid Daniel M, et al. Renal tubular PD-L1 (CD274) suppresses alloreactive human T-Cell responses. *Kidney Internl* 2010;78(1):38–47. <https://doi.org/10.1038/ki.2010.97>.
 92. Taguchi Atsuhiko, Kaku Yusuke, Ohmori Tomoko, Sharmin Sazia, Ogawa Minetaro, Sasaki Hiroshi, et al. Redefining the in vivo origin of metanephric nephron progenitors enables generation of complex kidney structures from pluripotent stem cells. *Cell Stem Cell* 2014;14(1):53–67. <https://doi.org/10.1016/j.stem.2013.11.010>.
 93. Takasato M, Er PX, Becroft M, Vanslambrouck JM, Stanley EG, Elefanty AG, Little MH. Directing human embryonic stem cell differentiation towards a renal lineage generates a self-organizing kidney. *Nature Cell Biol* 2014;16(1):118–26. <https://doi.org/10.1038/ncb2894>.
 94. Takasato Minoru, Er Pei X, Chiu Han S, Maier Barbara, Baillie Gregory J, Ferguson Charles, et al. Kidney organoids from human IPS cells contain multiple lineages and model human nephrogenesis. *Nature* 2015;526(7574):564–8. <https://doi.org/10.1038/nature15695>.

95. Tanigawa Shunsuke, Islam Mazharul, Sharmin Sazia, Naganuma Hidekazu, Yoshimura Yasuhiro, Haque Fahim, et al. Organoids from nephrotic disease-derived iPSCs identify impaired NEPHRIN localization and slit diaphragm formation in kidney podocytes. *Stem Cell Reports* 2018;11(3):727–40. <https://doi.org/10.1016/j.stemcr.2018.08.003>.
96. Tanigawa Shunsuke, Tanaka Etsuko, Miike Koichiro, Ohmori Tomoko, Inoue Daisuke, Cai Chen Leng, et al. Generation of the organotypic kidney structure by integrating pluripotent stem cell-derived renal stroma. *Nature Commun* 2022;13(1):1–15. <https://doi.org/10.1038/s41467-022-28226-7>.
97. Tekguc Murat, Wing James Badger, Osaki Motonao, Long Jia, Sakaguchi Shimon. Treg-Expressed CTLA-4 depletes CD80/CD86 by trogocytosis, releasing free PD-L1 on antigen-presenting cells. *Proceedings of the National Academy of Sciences of the United States of America* 2021;118(30):1–12. <https://doi.org/10.1073/pnas.2023739118>.
98. Tran Tracy, Lindström Nils O, Ransick Andrew, Brandine Guilherme De Sena, Guo Qiuyu, Kim Albert D, et al. In vivo developmental trajectories of human podocyte inform in vitro differentiation of pluripotent stem cell-derived podocytes. *Developmental Cell* 2019;50(1):102–16. <https://doi.org/10.1016/j.devcel.2019.06.001>.
99. Trautmann Agnes, Bodria Monica, Ozaltin Fatih, Gheisari Alaleh, Melk Anette, Azocar Marta, et al. Spectrum of steroid-resistant and congenital nephrotic syndrome in children: the podonet registry cohort. *Clin J Am Society of Nephrol* 2015;10(4):592–600. <https://doi.org/10.2215/CJN.06260614>.
100. Uchino Shigehiko, Kellum John A, Bellomo Rinaldo, Doig Gordon S, Morimatsu Hiroshi, Morgera Stanislaw, et al. Acute renal failure in critically ill patients: a multinational, multicenter study. *JAMA* 2005;294(7):813–8. <https://doi.org/10.1001/jama.294.7.813>.
101. Unbekandt Mathieu, Davies Jamie A. Dissociation of embryonic kidneys followed by reaggregation allows the formation of renal tissues. *Kidney Internat* 2010;77(5):407–16. <https://doi.org/10.1038/ki.2009.482>.
102. Vaidya Vishal S, Ozer Josef S, Dieterle Frank, Collings Fitz B, Ramirez Victoria, Troth Sean, et al. Kidney injury molecule-1 outperforms traditional biomarkers of kidney injury in preclinical biomarker qualification studies. *Nature Biotechnol* 2010;28(5):478–85. <https://doi.org/10.1038/nbt.1623>.
103. Wearn JT, Richards AN. Observations on the composition of glomerular urine, with particular reference to the problem of reabsorption in the renal tubules. *Am J Physiol-Legacy Content* 1924;71(1):209–27. <https://doi.org/10.1152/ajplegacy.1924.71.1.209>.
104. Wehmeier C, Hönger G, Cun H, Amico P, Hirt-Minkowski P, Georgalis A, et al. Donor specificity but not broadness of sensitization is associated with antibody-mediated rejection and graft loss in renal allograft recipients. *Am J Transplantation* 2017;17(8):2092–102. <https://doi.org/10.1111/ajt.14247>.
105. Whiting PH, Brown PAJ. The relationship between enzymuria and kidney enzyme activities in experimental gentamicin nephrotoxicity. *Renal Failure* 1996;18(6):899–909. <https://doi.org/10.3109/08860229609047716>.
106. Wiggins RC. The spectrum of podocytopathies: a unifying view of glomerular diseases. *Kidney Internat* 2007;71(12):1205–14. <https://doi.org/10.1038/sj.ki.5002222>.
107. Williams Robert C, Opelz Gerhard, McGarvey Chelsea J, Jennifer Weil E, Chakkera Harini A. The risk of transplant failure with HLA mismatch in first adult kidney allografts from deceased donors. *Transplantation* 2016;100(5):1094–102. <https://doi.org/10.1097/TP.0000000000001115>.
108. Wilson Sean B, Howden Sara E, Vanslambrouck Jessica M, Dorison Aude, Alquicira-Hernandez Jose, Powell Joseph E, Little Melissa H. DevKidCC allows for robust classification and direct comparisons of kidney organoid datasets. *Genome Med* 2022;14(1):1–25. <https://doi.org/10.1186/s13073-022-01023-z>.
109. Wing James B, Tekguc Murat, Sakaguchi Shimon. Control of germinal center responses by T-follicular regulatory cells. *Frontiers in Immunol* 2018;9(1910):1–12. <https://doi.org/10.3389/fimmu.2018.01910>.
110. Wu Haojia, Uchimura Kohei, Donnelly Erinn L, Kirita Yuhei, Morris Samantha A, Humphreys Benjamin D. Comparative analysis and refinement of human PSC-derived kidney organoid differentiation with single-cell transcriptomics. *Cell Stem Cell* 2018;23(6):869–81. <https://doi.org/10.1016/j.stem.2018.10.010> e8doi:.
111. Xia Yun, Nivet Emmanuel, Sancho-Martinez Ignacio, Gallegos Thomas, Suzuki Keiichiro, Okamura Daiji, et al. Directed differentiation of human pluripotent cells to ureteric bud kidney progenitor-like cells. *Nature Cell Biol* 2013;15(12):1507–15. <https://doi.org/10.1038/ncb2872>.
112. Xinaris Christodoulos, Benedetti Valentina, Rizzo Paola, Abbate Mauro, Corna Daniela, Azzollini Nadia, et al. In vivo maturation of functional renal organoids formed from embryonic cell suspensions. *J Am Society of Nephrol* 2012;23(11):1857–68. <https://doi.org/10.1681/ASN.2012050505>.
113. Xu Huaigeng, Wang Bo, Ono Miyuki, Kagita Akihiro, Fujii Kaho, Sasakawa Noriko, et al. Targeted disruption of HLA genes via CRISPR-Cas9 generates iPSCs with enhanced immune compatibility. *Cell Stem Cell* 2019;24(4):566–78. <https://doi.org/10.1016/j.stem.2019.02.005> e7.
114. Yang Yong Guang, Sykes Megan. Xenotransplantation: current status and a perspective on the future. *Nature Rev Immunol* 2007;7(7):519–31. <https://doi.org/10.1038/nri2099>.
115. Yokote Shinya, Matsunari Hitomi, Iwai Satomi, Yamanaka Shuichi, Uchikura Ayuko, Fujimoto Eisuke, et al. Urine excretion strategy for stem cell-generated embryonic kidneys. *Proceedings of the National Academy of Sciences of the United States of America* 2015;112(42):201507803. <https://doi.org/10.1073/pnas.1507803112>.
116. Zhang Shun, Wan Zhengpeng, Kamm Roger D. Vascularized organoids on a chip: strategies for engineering organoids with functional vasculature. *Lab on a Chip* 2021;21(3):473–88. <https://doi.org/10.1039/d0lc01186j>.
117. Zhang Yu Shrike, Arneri Andrea, Bersini Simone, Shin Su-Ryon, Zhu Kai, Goli-Malekabadi Zahra, et al. Bioprinting 3D microfibrous scaffolds for engineering endothelialized myocardium and heart-on-a-chip. *Biomaterials* 2016;110:45–59. <https://doi.org/10.1016/j.biomaterials.2016.09.003> December.